

mass concentrations. The size of emitted particles is very strongly related to the temperature of the wood sample and composition of the atmosphere.

Acknowledgement

Presented results were obtained within the project NETME CENTRE PLUS (LO1202), created with financial support from the Ministry of Education, Youth and Sports of the Czech Republic under the „National Sustainability Programme I”.

References:

- [1] HINDS, William C. Aerosol technology: properties, behavior, and measurement of airborne particles. 2nd ed. New York: Wiley, c1999. ISBN 978-0-471-19410-1.
- [2] FARA, M. Toxikologicky závažné látky v emisích PM10, [online]. Praha, [cit. 20121219]. Projekt. ČHMU. 2003 Available at: http://old.chmi.cz/uoco/prj/vav_740_3_02/dp1d od.pdf
- [3] KURFÜRST, Jiří, ed. Kompendium ochrany kvality ovzduší. Chrudim: *Vodní zdroje Ekomonitor*, 2008. ISBN 978-80-86832-38-8.
- [4] Kroschwitz J. and Seidel A., Kirk-Othmer encyclopedia of chemical technology, 5th ed. Hoboken, N.J, Wiley-Interscience, c2007, p. 26
- [5] Cachim P. and Franssen J., Comparison between the charring rate model and the conductive model of Eurocode 5, *Fire and OBAIDULLAH, M., et al. A review on particle emissions from small scale biomass combustion. International Journal of Renewable Energy Research (IJRER)*, 2012, 2.1: 147-159.
- [6] TISSARI, Jarkko, et al. A novel field measurement method for determining fine particle and gas emissions from residential wood combustion. *Atmospheric Environment*, 2007, 41.37: 8330-8344.
- [7] SIPPULA, O. Fine particle formation and emission in biomass combustion. Ph D thesis. Department of Environmental Science. University of Eastern Finland. Finland. 2010. ISBN 978-952-5822-14-4.
- [8] OBERNBERGER, I., BRUNNER, T., BARNTHALER, G. Fine particle emissions from Modern Austrian small scale biomass combustion plants. *15th European biomass conference and exhibition*. 7-11 May 2007, Germany:1546-57.