































**Table 1. Identification of Microalgae in Ponds in Pekanbaru City.**

No	Classification	Observations under a Light Microscope	Image morphology cell sourced from reading material	Description of microalgae
1	Kingdom : Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Chlorococcales Family : Scenedesmaceae Genus : Scenedesmus Species : <i>Scenedesmus quadricauda</i>			<ul style="list-style-type: none"> <li>- Unicellular Cell wall with hemicellulosic and sporopolleninic layers</li> <li>- Cells unite and connect.</li> <li>- Can to move</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
2	Kingdom : Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Chlorococcales Family : Scenedesmaceae Genus : Scenedesmus Species : <i>Scenedesmus obliquus</i>			<ul style="list-style-type: none"> <li>- Form flagella isocontae</li> <li>- Bright green</li> <li>- Live in the colony</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
3	Kingdom : Plantae Phylum : Chlorophyta Class : Chlorophyceae Order: Chlorococcales Family : Scenedesmaceae Genus : Scenedesmus Species : <i>Scenedesmus dimorphus</i>			<ul style="list-style-type: none"> <li>- Colonize</li> <li>- Cells are elliptical to lanceolate</li> <li>- Thorn</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
4	Kingdom : Plantae Phylum : Chlorophyta Class: Chlorophyceae Order : Chlorococcales Family : Scenedesmaceae Genus : Scenedesmus Species : <i>Scenedesmus acuminatus</i>			<ul style="list-style-type: none"> <li>- Shaped</li> <li>- Spiky</li> <li>- Green color</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	

5	Kingdom :Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Sphaeropleales Family : Hydrodictyaceae Genus : Pediasstrum Species: <i>Pediasstrum biradiatum</i>			<ul style="list-style-type: none"> <li>- Reproduce asexually by producing autocolonies</li> <li>- The protoplast of each parent cell gives rise to zoospores.</li> <li>- Number of colonized cells</li> <li>- Disc-shaped and marked like a peripheral horn</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
6	Kingdom :Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Sphaeropleales Family : Hydrodictyaceae Genus : Pediasstrum Species: <i>Pediasstrum duplex may</i>			<ul style="list-style-type: none"> <li>- The flagella are always the same length, called isocontae</li> <li>- Consists of the pigment Chlorophyta A</li> <li>- Food reserves are carbohydrates in the form of flour and protein in the form of pyrenoids</li> <li>- Cell wall is composed of cellulose</li> <li>- It is an order of colonizing Chlorococcales</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
7	Kingdom :Plantae Phylum : Chlorophyta Class : Chlorophyceae Order: Sphaeropleales Family : Hydrodictyaceae Genus : Pediasstrum Species : <i>Pediasstrum duplex var</i>			<ul style="list-style-type: none"> <li>- The body is flat and resistant to drought</li> <li>- Spores form hypospores</li> <li>- Cells are green, red, and brown</li> <li>- Small in size, unicellular, and have flagella</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
8	Kingdom : Plantae Phylum : Charophyta Class : Chlorophyceae Order : Desimidiales Family : Closteriaceae Genus : Closterium Species : <i>Closterium parvulum</i>			<ul style="list-style-type: none"> <li>- Shaped like an elongated sickle</li> <li>- Curved and tapered at the ends</li> <li>- There have chloroplasts so they can photosynthesize</li> <li>- They has many vacuoles at the ends.</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
9	Kingdom : Chromista Phylum : Chlorophyta Class : Chlorophyceae Order : Chlorococcales Family : Chlorellacea Genus : Chlorella Species: <i>Chlorella</i> sp.			<ul style="list-style-type: none"> <li>- The cells are spherical in shape and small in size</li> <li>- There have cell walls</li> <li>- This algae has chlorophyll</li> <li>- Includes green algae</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	

10	Kingdom : Eubacteria Phylum : Cyanobacteria Class : Cyanophyceae Order : Spirulinales Family : Spirulinaceae Genus : Spirulina Species : <i>Spirulina</i> sp.			<ul style="list-style-type: none"> <li>- The cells are thread-shaped colonies</li> <li>- The threads are unbranched</li> <li>- Cells are like green beads</li> <li>- Between the cells of the body there are empty cells</li> <li>- Includes green algae</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
11	Kingdom : Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Sphaeropleales Family : Scenedesmaceae Genus : Coelastrum Species : <i>Coelastrum microforum</i>			<ul style="list-style-type: none"> <li>- The cell walls of these organisms are double layered.</li> <li>- This wall layer contains many short projections called papillae.</li> <li>- Binds coenobium cells together with a dividing nucleus.</li> <li>- A vegetative cell exhibits a peripheral disposition and contains a single located ribosomal nucleolus.</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
12	Kingdom : Eubacteria Phylum : Cyanobacteria Class : Cyanophyceae Order : Chroococcales Family : Microcystaceae Genus : Microcystis Species : <i>Microcystis aeruginosa</i>			<ul style="list-style-type: none"> <li>- They have small cells</li> <li>- They does not have individual membranes</li> <li>- The color of the protoplast looks dark or brown</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	
13	Kingdom : Plantae Phylum : Chlorophyta Class : Chlorophyceae Order : Sphaeropleales Family : Scenedesmaceae Genus : Tetrastrum Species : <i>Tetrastrum Heteracarthum</i>			<ul style="list-style-type: none"> <li>- Live in colonies</li> <li>- They have branches</li> <li>- Divide cells</li> </ul>
		Magnification (100x)	Source: Yunfang (1995)	

**Information:**

- Personal documentation
- Identification using the Yunfang (1995) manual
- Image source image from Algae Base ([www.algaebase.org](http://www.algaebase.org))

decomposition rate of saprist. Another physical property of peat soil is irreversible drying. Dried peat, with a moisture content of <100% (by weight), cannot absorb any more water when wet [4]. This dried peat has the same properties as dry wood, which is easily washed away by the flow of water and is flammable.




Figure 1. Average abundance of microalgae in fish farming soil ponds in Pekanbaru Cit

Peatlands in Pekanbaru City generally have a high level of acidity with a pH range of 3-5. Oligotrophic peat that has a quartz sand substrate in Berengbengkel, Central Kalimantan has a pH range of 3.25 – 3.75 [5]. Meanwhile, the peat around Air Sugihan Kiri, South Sumatra has a higher pH range between 4.1 to 4.3 [2]. Peatlands naturally have a low fertility level, characterized by pH, and low content of micro and macronutrients. Based on the level of fertility Fresher [6] divides peat into three types, namely Eutrophic (fertile), mesotrophic (moderate), and oligotrophic (infertile). Topogenous peat that is shallow and influenced by groundwater and rivers is generally categorized as mesotrophic to eutrophic, while embryogenic peat that is only affected by rainwater is categorized as oligotrophic [7].

From table 1, the types of microalgae were successfully described and classified during the study. The class Chlorophyceae is a group that is commonly found in the waters of fish farming soil ponds in the city of Pekanbaru. Several previous studies [8] stated that Chlorophyceae is the most abundant group of microalgae living in general waters in Riau Province. Microalgae are microscopic unicellular algae, usually found in freshwater and marine environments and exist individually or in chains or groups [9] – [11]. Microalgae are a very diverse group of organisms, including both prokaryotes and eukaryotes, and spanning 14 phyla with described examples from

almost every possible habitat. It is estimated that 200,000 – 800,000 species of microalgae are found widely, of which only about 35,000. Microalgae habitats include freshwater, seawater, soil, and extreme environments [12]–[14].

Based on the highest number of microalgae species found during the study in week 1, week 2, and week 3, it showed that the highest number of species were *Spirulina plantensis* (316440 Cell/L), *Chlorella vulgaris* (38748 Cell/L), and *Scenedesmus quadricauda* (31553 Cell/L). (Figure 3). The types of microalgae found in the waters of this fish farming pond are different from the types of microalgae found by [15], [16], in the waters of the islands of Gumilamo and Magaliho, North Halmahera. He reported that his study found seven genera of diatoms, namely: *Chaetoceros* sp. with the highest abundance (5,061 cells/L), followed by *Nitzschia* sp. (611 cells/L), *Thalassionema* sp. (569 cells/L), *Skeletonema* sp. (446 cells/L), *Coscinodiscus* sp. (176 cells/L), *Navicula* sp. (40 cells/L) and *Amphora* sp. (11 cells/L).

Based on the results of the analysis of the abundance and dominance of microalgae conducted by previous open researchers [17]–[21], several types of microalgae are abundant and have the potential to be developed as natural feed on a larger scale or industry, one of which is microalgae as a substitute (substitution) sources of carbohydrates and animal protein in artificial feeds for cultured fish. Experts [22]–[26] also stated that based on existing technology, and research that continues to be developed today, the species or genus *Chlorella* and *Spirulina* are the most likely types of microalgae for development towards mass cultivation.

However, the use of microalgae as an artificial feed substitute for fish, to support aquaculture activities requires several suitability criteria for the cultivation of the developer, especially the feasibility of the quality of the location or waters in the location to be developed [27]–[30]. Some of the eligibility criteria include the results of direct measurements of the abundance of a type of microalgae in nature, eligibility criteria for physical and chemical properties of soil or water at the location or field, community support, land topography, and supporting facilities [31]–[34]. In the results of this study, almost all of the eligibility criteria have been answered.

Based on Figure 2 above, the highest density of microalgae is, *Spirulina platensis* 33%, *Chlorella Vulgaris* 29%, and *Scenedesmus quadricauda* 10%.




Figure 2. The density of Microalgae from Groundwater Pond Fish Cultivation in Pekanbaru City




Figure 3. Biomass of the three most abundant types of microalgae from fish farming soil ponds in Pekanbaru City which are cultured on a laboratory scale

The diagram above shows that Scenedesmus quadricauda is higher than Spirulina platensis and Chlorella Vulgaris. Meanwhile, the research reports that the microalgae found in the waters of the river and estuary of the Pelangan River consist of 2 divisions, namely the Chrysophyta division and the Chlorophyta division. The total types of microalgae found at all sampling points were 85 species, including 13 species at the point I, 23 species at point II, 7 species at point III, 24 species at point IV, and 18 species at point V. The highest microalgae species diversity index ( $H^1=3,010$ ) was found at sampling point IV, medium category, and also with the highest number of species (24 species) with 77 individuals. The highest total abundance (1,540 ind/L) occurred at sampling point II. While the type of microalgae with the highest abundance was Chlorococcum humicola: 1045 ind/L occurred at sampling point II. At sampling points, I, II, and III based on importance, the dominant species were Chlorococcum humicola and Coscinodiscus lacustris. While at points IV and V, the dominant species was Thalassiotrix fruenfeldii with a dominance percentage of 15.795% and 21.840%, respectively.

Based on the results of the calculation of the highest dry weight biomass of microalgae found in pond waters of fish culture in Pekanbaru City is Scenedesmus Quadricauda 0.26 (gr). The 3 highest types of biomass found were Scenedesmus Quadricauda 0.26 (gr), Spirulina platensis 0.13 (gr), and Chlorella vulgaris 0.11 (gr). According to experts [28], [35]–[39], microalgae are known to have superior characteristics as potential raw materials for commercial biodiesel production. However, it is still hit by the high cost of production when compared to fossil-based diesel fuel. Therefore, various efforts to improve its economic feasibility must be carried out. One of the surefire steps in solving this problem is to maximize the production of microalgae biomass, so that lipid productivity and biodiesel productivity can be increased. Optimization of microalgae biomass production was carried out on four main cultivation parameters, such as supplementation of carbon dioxide (CO<sub>2</sub>), composition of growth media, optimization of environmental conditions and addition of growth factors/hormones [40]–[44]. Operational microalgae cultivation under optimal conditions will maximize its biomass production until it can finally achieve maximum biodiesel productivity [45]–[51].

Table 2. Results of Water Quality Measurement in Ponds in Pekanbaru City.

Water Quality	Value range
Temperature (°C)	28,5-29,33
Turbidity (cm)	18,25-38
pH	6,00
DO (mg/L)	2,45-6,53
CO <sub>2</sub> Bebas (mg/L)	3,00-14,98
Nitrat (NO <sub>3</sub> -)	0,1-0,3
Fosfat (PO <sub>4</sub> <sup>3-</sup> )	0,005-0,008

Based on research reports from other researchers [25],[34],[41],[54], microalgae are photosynthetic microorganisms that utilize carbon dioxide and sunlight to form biomass and produce about 50 percent of the oxygen in the atmosphere. There are four types of microalgae, namely Bacillariophyceae (diatoms), Chlorophyceae (green algae), Chrysophyceae, (gold algae) and Cyanophyceae (blue algae) [52], [53]. Although Indonesia has a high diversity of microalgae, its potential is still not fully utilized. In many countries, microalgae have been used as agents for biofuel production, because microalgae are capable of producing high levels of fatty acids and carbohydrates. Through the

esterification process, microalgae fatty acids can be converted into biodiesel.

The potential of microalgae is very promising to be used as an agent in the production of biofuels in the future because it has advantages in its growth, namely fast, high productivity, does not require a large area of land for breeding, and can use water for growing nutrients [54]. In addition, the use of microalgae as a source of biodiesel does not interfere with the food supply. This is because microalgae do not compete with foodstuffs [35]. Another potential of using microalgae is its ability to use carbon dioxide and convert it into oxygen. This ability can create a clean environment from CO<sub>2</sub> gas pollutants so that it can suppress the effects of global warming.

*Chlorella Vulgaris* is a type of microalgae that is commonly found in Indonesian waters, but its potential for producing biodiesel has not been developed [55]. A local strain of *Chlorella Vulgaris* has been developed for biodiesel production. Microalgae *Chlorella Vulgaris* was grown in DG11 medium with a CO<sub>2</sub> supply. Optimization of media and fermentation conditions was carried out so that the fermentation could produce biomass with high fatty acid content [56]. The biomass content of microalgae produced in the fermentation process is determined by measuring its optical density, while the content and composition of fatty acids in biomass will be determined by titration and GC-MS methods.

Microalgae *Chlorella Vulgaris* can be grown well on BG-11 media which contains several mineral mixtures. The growth of this microalgae takes place more optimally when a cell starter with a concentration of 16% (v/v) is used. The local strain of *Chlorella Vulgaris* biomass can produce lipids with quite high levels, namely 31% (v/v) when extracted with a mixed solvent of n-hexane-ethanol (1:1). Analysis by GC-MS can be seen that the lipid content in *Chlorella Vulgaris* is composed of dominant fatty acids, namely pentadecanoic acid, hexadecanoic acid, heptadecanoic acid, stearic acid and nonadecanoic acid [22], [57]. Biodiesel production from the microalga *Chlorella Vulgaris* has been carried out both ex-situ and in-situ. The in-situ process of lipid transesterification in the microalgae *Chlorella Vulgaris* biomass can produce higher biodiesel yields than ex-situ [58], [59].

In-situ biodiesel production can take place optimally when subjected to ultrasonication at a power of 25 kHz (270W), using n-hexane co-solvent, and with a ratio of biomass to methanol weight of 3:50. The comparison of 3:50 resulted in a higher percentage of biodiesel conversion and

biodiesel yield of 20.31% w/w and 3.87% w/w using the sonication method. This in situ process can produce biodiesel of 20.31% w/w. These results can prove that microalgae that are abundant in Indonesian waters can be developed as a source of biodiesel production so that it can help supply alternative energy to fossil fuels. The scale of biodiesel production by pilot plant and industry needs to be supported by the government, either by policy or by providing adequate infrastructure, given its enormous potential.

## 4 Conclusion

From this research, 13 species of microalgae have been found that live in fish culture pond habitat in Pekanbaru City, which consist of *Scenedesmus quadricauda*, *Pediastrum duplex meyeri*, *Tetrastrum heteracanthum*, *Spirulina plantensis*, *Microcystis aeruginosa*, *Chlorella vulgaris*, *Scenedesmus dimorphus*, *Pediastrum duplex var.*, *Closterium parvulum*, *Coelastrum microforum*, *Scenedesmus acuminatus*, *Pediastrum biradiatum* and *Scenedesmus obligus*. The most common type was found in *Spirulina platensis* (316440 cells/L), and the least common was *Microcystis aeruginosa* (1228 cells/L). The highest density was found in *Spirulina platensis* 33 %, and the lowest density was in *Microcystis aeruginosa* 1 %. Meanwhile, the order of biomass from the highest to the lowest of the 3 most common types from ground pool waters was *Jenensi Scenedesmus quadricauda* 0.26 (gr), *Spirulina platensis* 0.13 (gr), and *Chlorella vulgaris* 0.11 (gr). Then the water quality parameters recorded in the soil pond include temperature 29°C, pH 6, brightness 21 Cm, dissolved oxygen 3.81 Mg/L, CO<sub>2</sub> 5.75, Mg/L, Phosphate 0.55 mg/L, and Nitrate 1, 26 mg/L..

Conflict of interest: None.

### References:

- [1] D. Darjono, "Experience of Law Enforcement Relating to Fires in Plantation Areas and HTI Swamp Peat," in *Fires in swamp/peat land in Sumatra: problems and solutions*. Seminar Proceedings, Palembang, South Sumatra, 10 – 11 December 2003, no. 19, 2004, pp. 73–104. doi: 10.17528/cifor/001568.
- [2] T. Van Ty et al., "Scenario-based impact assessment of land use/cover and climate changes on water resources and demand: a case study in the Srepok Basin, Vietnam -



- Cambodia,” *Water Resource Manag.*, vol. 26, no. 5, pp. 1387–1407, 2012,
- [3] S. Chittapun, P. Pholpunthin, and H. Segers, “Contribution to the knowledge of Thai microfauna diversity: Notes on rare peat swamp Rotifera, with the description of a new *Lecane Nitzsch*, 1872,” *Hydrobiologia*, vol. 501, pp. 7–12, 2003
- [4] P. A. Abrams et al., “Eradication of feral pigs from Australia,” *Conserv. Biol.*, vol. 10, no. 1, pp. 1–11, 2001.
- [5] S. Cui, Q. Zhou, and L. Chao, “Potential hyperaccumulation of Pb, Zn, Cu and Cd in enduring plants distributed in an old smeltery, northeast China,” *Environ. Geol.*, vol. 51, no. 6, pp. 1043–1048, 2007
- [6] D. P. Thakur and C. K. Lin, “Water quality and nutrient budget in closed shrimp (*Penaeus monodon*) culture systems,” *Aquac. Eng.*, 2003
- [7] J. Markiewicz-Patkowska, A. Hursthouse, and H. Przybyla-Kij, “The interaction of heavy metals with urban soils: Sorption behavior of Cd, Cu, Cr, Pb and Zn with a typical mixed brownfield deposit,” *environment. Int.*, vol. 31, no. 4, pp. 513–521, 2005
- [8] P. Arulmurugan, S. Nagaraj, and N. Anand, “Biodiversity of fresh water algae from Guindy campus of Chennai, India,” *J. Ecobiotechnology*, vol. 3, no. 10, pp. 19–29, 2011, [Online]. Available: <http://journal-ecobiotechnology.com/>
- [9] D. Remias, U. Karsten, C. Lütz, and T. Leya, “Physiological and morphological processes in the Alpine snow alga *Chloromonas nivalis* (Chlorophyceae) during cyst formation,” *Protoplasma*, 2010
- [10] C. Rosenberg and J. Ramus, “Ecological growth strategies in the seaweeds *Gracilaria foliifera* (Rhodophyceae) and *Ulva* sp. (Chlorophyceae): Soluble nitrogen and reserve carbohydrates,” *Mar. Biol.*, 1982
- [11] C. Wang et al., “How DNA barcoding can be more effective in microalgae identification: A case of cryptic diversity revelation in *Scenedesmus* (Chlorophyceae),” *Sci. Rep.*, vol. 6, 2016
- [12] L. Rhodes et al., “Cryopreservation of economically valuable marine micro-algae in the classes Bacillariophyceae, Chlorophyceae, Cyanophyceae, Dinophyceae, Haptophyceae, Prasinophyceae, and Rhodophyceae,” *Cryobiology*, vol. 52, no. 1, pp. 152–156, 2006
- [13] O. Gómez-Espinoza, M. Guerrero-Barrantes, K. Meneses-Montero, and K. Núñez-Montero, “Identification of a microalgae collection isolated from Costa Rica by 18s rDNA sequencing,” *Acta Biologica Colombiana*, vol. 23, no. 2, 2018.
- [14] M. T. Jahn, K. Schmidt, and T. Mock, “A novel cost effective and high-throughput isolation and identification method for marine microalgae,” *Plant Methods*, vol. 10, no. 1, 2014
- [15] D. Nlawati, “Preliminary Study of Biodiesel Synthesis from Lipid Microalgae *Chlorella vulgaris* Based on Walne Medium Through Esterification and Transesterification Reaction. Book: Thesis Univ. Indonesia., 2012.
- [16] R. Nasrul Sani, F. Choirun Nisa, R. Dewi Andriani, and J. Mahar Maligan, “Analysis of Yield and Phytochemical Screening of Microalgae Extracts-Sani, et al,” 2014.
- [17] M. Daroch, S. Geng, and G. Wang, “Recent advances in liquid biofuel production from algal feedstocks,” *Applied Energy*, vol. 102, pp. 1371–1381, 2013.
- [18] K. Menzel and A. Wild, “Fatty Acid Composition in the Lipids of some Marine Chlorococcales and Eustigmatales,” *Zeitschrift fur Naturforsch. - Sect. C J. Biosci.*, 1989.
- [19] A. Neori et al., “Integrated aquaculture: Rationale, evolution and state of the art emphasizing seaweed biofiltration in modern mariculture,” *Aquaculture*. 2004.
- [20] S. Sathya and S. Srisudha, “Isolation and Identification of Freshwater Microalgal Strains-Potential for Biofuel Production,” *Int. J. Recent Sci. Res.*, vol. 4, no. August, 2018.
- [21] R. Pate, G. Klise, and B. Wu, “Resource demand implications for US algae biofuels production scale-up,” *Appl. Energy*, vol. 88, no. 10, pp. 3377–3388, 2011.
- [22] G. De Bhowmick, G. Subramanian, S. Mishra, and R. Sen, “Raceway pond cultivation of a marine microalgae of Indian origin for biomass and lipid production: A case study,” *Algal Res.*, vol. 6, no. PB, pp. 201–209, 2014
- [23] U. (Sweden). D. of F. S. Pickova, Jana (Swedish University of Agricultural Sciences), U. (Sweden). D. of F. S. Mraz, Jan (Swedish University of Agricultural Sciences, and U. (Sweden). D. of F. S. Zajic, Tomas (Swedish University of Agricultural Sciences, “Common carp and aquaculture,” 2013.
- [24] T. C. O’Keefe et al., “Editorial,” *Ecology*, vol. 48, no. 1, pp. 1–13, 2014.
- [25] M. Jaya Prakash Goud, D. Seshikala, and M. A. Singara Charya, “Antibacterial activity and biomolecular composition of certain fresh water micro-algae from River Godavari

- (India),” *Sci. World J.*, vol. 2, no. 3, pp. 19–23, 2007.
- [26] S. Chinnasamy, A. Bhatnagar, R. W. Hunt, and K. C. Das, “Microalgae cultivation in a wastewater dominated by carpet mill effluents for biofuel applications,” *Bioresour. Technol.*, 2010.
- [27] K. J. Patil, H. Nandre, R. T. Mahajan, C. Banerjee, P. Shukla, and R. Bandopadhyay, “Plant growth stimulating activity of fresh water micro algae,” *Int. J. Adv. res. Journalwww.journalijar.com Int. J. Adv. Res.*, vol. 3, no. 5, pp. 284–288, 2015.
- [28] A. Varithaa, A. Dharmaa, E. Munafa, N. Nasir, and Afrizal, “Isolation oil producing micro algae *Chlamydomonas snowii* from tropical fresh water, Indonesia,” *Res. J. Pharm. Biol. Chem. Sci.*, vol. 4, no. 4, pp. 1462–1470, 2013.
- [29] P. Coltelli, L. Barsanti, V. Evangelista, A. M. Frassanito, and P. Gualtieri, “Water monitoring: Automated and real time identification and classification of algae using digital microscopy,” *Environ. science. Process. Impacts*, vol. 16, no. 11, 2014.
- [30] U. Sabeela Beevi and R. K. Sukumaran, “Cultivation of the fresh water microalga *Chlorococcum* sp. RAP13 in sea water for producing oil suitable for biodiesel,” *J. Appl. Phycol.*, vol. 27, no. 1, pp. 141–147, 2014.
- [31] [31] E. Juwita, K. Soewardi, and Y. Yonvitner, “Habitat Conditions and Mangrove Ecosystem in Simpang Pesak District, East Belitung for Development of Shrimp Pond,” *J. Mns. and Lingkungan*, 2015, doi: 10.22146/jml.18725.
- [32] E. Nunez-Lara and J. E. Arias-Gonzales, “The relationship between reef fish community structure and environmental variables in the southern Mexican Caribbean,” *J. Fish Biol.*, vol. 53, no. supp. A, pp. 209–221, 1998,
- [33] E. Nunez-Lara and E. Arias Gonzalez, “The relationship between reef fish community structure and environmental variables in the southern Mexican Caribbean,” *J. Fish Biol.*, vol. 53, no. SUPPL. A, pp. 209–221, 1998
- [34] E. Núñez-Lara and J. E. Arias-González, “The relationship between reef fish community structure and environmental variables in the southern Mexican Caribbean,” *J. Fish Biol.*, vol. 53, no. sa, pp. 209–221, 1998, doi:10.1111/j.1095-8649.1998.tb01028.x.
- [35] G. Kanaya, S. Takagi, and E. Kikuchi, “Spatial dietary variations in *Laternula marilina* (Bivalva) and *Hediste* spp. (Polychaeta) along environmental gradients in two brackish lagoons,” *Mar. Ecol. prog. Ser.*, vol. 359, pp. 133–144, 2008.
- [36] P. Soudant et al., “The use of lipid emulsions for sterol supplementation of spat of the Pacific oyster, *Crassostrea gigas*,” *Aquaculture*, 2000
- [37] K. Matsumoto et al., “Effect of acute and chronic ammonia and nitrite exposure on oxygen consumption and growth of juvenile big bellied seahorse,” *Aquaculture*, vol. 7, no. 1, pp. 1–7, 2011
- [38] D. H. Lee, “Algal biodiesel economy and competition among bio-fuels,” *Bioresour. Technol.*, vol. 102, no. 1, pp. 43–49, 2011.
- [39] M. . Soheili and K. . Khosravi-Darani, “The potential health benefits of algae and micro algae in medicine: A review on *Spirulina platensis*,” *Curr. Nutrient. Food Science*, vol. 7, no. 4, pp. 279–285, 2011.
- [40] F. Alam, S. Mobin, and H. Chowdhury, “Third generation biofuel from Algae,” in *Procedia Engineering*, 2015, vol. 105, pp. 763–768.
- [41] R. Taylor and R. L. Fletcher, “Cryopreservation of eukaryotic algae - A review of methodologies,” *Journal of Applied Phycology*, vol. 10, no. 5. pp. 481–501, 1998.
- [42] B. H. Um and Y. S. Kim, “Review: A chance for Korea to advance algal-biodiesel technology,” *Journal of Industrial and Engineering Chemistry*, vol. 15, no. 1. pp. 1–7, 2009.
- [43] P. Coutteau and P. Sorgeloos, “The use of algal substitutes and the requirement for live algae in the hatchery and nursery rearing of bivalve molluscs: an international survey,” *J. Shellfish Research*, vol. 11. pp. 467–476, 1992.
- [44] R. Septory and B. Triyatmo, “The utilization of aquacultures wastewater as nutrient sources in *Nannochloropsis oculata* cultivation to prevent waters contamination in coastal area,” 2016.
- [45] K. Kumar, S. K. Mishra, A. Shrivastav, M. S. Park, and J. W. Yang, “Recent trends in the mass cultivation of algae in raceway ponds,” *Renewable and Sustainable Energy Reviews*. 2015.
- [46] Hadiyanto, M. Christwardana, and D. Soetrisnanto, “Phytoremediations of Palm Oil Mill Effluent (POME) by using aquatic plants and microalga for biomass production,” *J. Environ. science. Technol.*, vol. 6, no. 2, pp. 79–90, 2013.
- [47] K. E. Arthur, S. Kelez, T. Larsen, C. A. Choy, and B. N. Popp, “Tracing the biosynthetic source of essential amino acids in marine turtles using delta C-13 fingerprints,” *Ecology*, vol. 95, no. 5, pp. 1285–1293, 2014.

- [48] J. L. Blatti and M. D. Burkart, "Releasing stored solar energy within pond scum: Biodiesel from algal lipids," *J. Chem. Educ.*, vol. 89, no. 2, pp. 239–242, 2012.
- [49] P. Varshney, P. Mikulic, A. Vonshak, J. Beardall, and P. P. Wangikar, "Extremophilic micro-algae and their potential contribution in biotechnology," *Bioresource Technology*, vol. 184, pp. 363–372, 2015.
- [50] H. Coelho, T. Lopes da Silva, A. Reis, H. Queiroga, J. Serôdio, and R. Calado, "Fatty acid profiles indicate the habitat of mud snails *Hydrobia ulvae* within the same estuary: Mudflats vs. seagrass meadows," *Estuar. Coasts. Shelf Science*, vol. 92, no. 1, pp. 181–187, 2011.
- [51] J. L. Ruesink, "Epiphyte load and seagrass performance are decoupled in an estuary with low eutrophication risk," *J. Exp. March. Bio. Ecol.*, vol. 481, pp. 1–8, 2016
- [52] E. Kojima and K. Zhang, "Growth and hydrocarbon production of microalgae *Botryococcus braunii* in bubble column photobioreactors," *J. Biosci. Bioeng.*, vol. 87, no. 6, pp. 811–815, 1999.
- [53] M. D. Macías-Sánchez et al., "Biodiesel production from wet microalgal biomass by direct transesterification," *Fuel*, vol. 150, pp. 14–20, 2015.
- [54] B. R. Tampangallo, H. Herlinah, and M. C. Undu, "Incidence and Prevalence of WHITE SPOT SYNDROME VIRUS Infection in Plankton in Shrimp Farming Ponds," *J. Ris. Aquaculture*, 2018.
- [55] K. L. Straub et al., "The Vent and Seep Biota," *Nature*, vol. 4, no. 1, pp. 1–10, 2008.
- [56] M. Elmoraghy and I. Farag, "Bio-jet Fuel from Microalgae: Reducing Water and Energy Requirements for Algae Growth," *Int. J. Eng. Sci.*, vol. 1, no. 2, pp. 22–30, 2012
- [57] D. M. M. Dantas, A. R. F. Drummond, L. B. G. dos Santos, F. K. dos Santos, R. de S. Bezerra, and A. O. Gálvez, "Extração de óleo de microalga utilizando ultrassom com diferentes solventes visando a produção de biodiesel 1. 2010, pp. 1900–1904.
- [58] R. Praveenkumar, K. Shameera, G. Mahalakshmi, M. A. Akbarsha, and N. Thajuddin, "Influence of nutrient deprivations on lipid accumulation in a dominant indigenous microalgae *Chlorella* sp., BUM11008: Evaluation for biodiesel production," *Biomass and biodiesel production Bioenergy*, vol. 37, pp. 60–66, 2012.
- [59] [59] N. Mallick, S. Mandal, A. K. Singh, M. Bishai, and A. Dash, "Green microalgae *Chlorella vulgaris* as a potential feedstock for biodiesel," *J. Chem. Technol. Biotechnol.*, vol. 87, no. 1, pp. 137–145, 2012, doi:10.1002/jctb.2694.